

# What is StatPearls?

StatPearls provides exam review and selftesting resources.

#### Subscribed packages:

All Medical Specialties\*\*

Dental Package\*

National Board Dental Hygienist

Examination (NBDHE)\*

Nurse Practitioner\*\*

Nursing Certification\*\*

Nursing NCLEX (RN and PN)\*\*

Pharmacy\*\*

Physician Assistant\*\*

USMLE Steps 1, 2, and 3\*\*


## How do you get to StatPearls?

Statpearls can be found in the list of Databases.

You must register first through the LSUHSC-New Orleans registration portal using this link: LSUHSC-New Orleans Registration Portal

Once you are registered, you can log in at the StatPearls website


https://www.lsuhsc.edu/library/databases/statpearls.aspx



Q-Bank, Practice Test and CME/CE Finder

# Navigating Practioners I gating State elect earls State elect elec

**Get Started** 



Billing Info

Help & FAQs






Institutions


#### User Dash

The User Dash Is a quick overview of your account, focusing on upcoming quizzes and assignments.

It also includes a search that links to StatPearls overviews of various topics.

Below, you can find the selftest and continuing education resources.





## Take a Quiz

- Click on "Assigned Quizzes" in the "Assignments" group to access your quizzes.
- Uncompleted assignments appear in red.
- Completed assignments will show a score and possibly the option to review results.

#### Quiz: demonstration quiz 10.1.25

An otherwise healthy 23-year-old woman presents after a motor vehicle collision. A cervical collar is in place and oxygen is being delivered via a non-rebreather mask. According to the paramedics who transported her to the emergency department, the patient was recently conversing, but now she is unconscious, has a Glasgow Coma Scale score of 8, and requires intubation. Which of the following signs would most strongly preclude the use of nasotracheal intubation for this patient?

- O A. Scalp laceration
- B. Hypotension and tachycardia
- C. Flail chest
- D. Periorbital ecchymosis

Adult Lab Normals 7 Pediatric Lab Normals 7

Question 1 of 10

Send Comment to Editors

Quit and Exit Quiz

Next Question >

#### **Assignment Details**

Your Test Duration: 0.0.127

Total Allowed Duration: Unlimited

End Date and Time: 10/2/2025 9:38:00 AM

#### Question Details

Last Updated: 3/14/2024 2:23:27 PM

Quiz Type: Test Mode

Question Author: Thomas B. Folino George Mckean

Article/Question Editor: Lance J. Parks

#### Tips for Quizzes:

Click on the text to mark out options before selecting your answer.

For untimed quizzes, you may exit and resume, but timed quizzes will count time from start of the quiz and won't allow you to resume once the time has run out.

**NOTE:** The timer will reset for every new question. Use an external timer to track elapsed time.

#### **Out of Time?**

Sorry, this is a timed assignment with an allowed time of 1 min. You have started the assignment and your current total assignment time is 33 min. Please contact your instructor for any extension requests or to reset your assignment to be taken again.

Back

If you run out of time during a quiz or try to complete it after the due date, you will see an error message like the one above. Please reach out to your instructor about extensions or re-takes (or if they may have set the wrong due date on the assignment).

# Viewing Results

If your instructor has allowed students to view results, you can see the questions you missed or failed to answer on assignments, as well as the correct answer and study resources.

Otherwise, you'll just see a percent right/wrong.

A 17-year-old boy presents with an empty right hemiscrotum. The left testicle is normal. He is healthy, and a right testicle is seen in the right paracolic gutter. Which of the following is the most appropriate management? 2. Start luteinizing hormone-releasing hormone. 3. Reassure the patient. 4. Perform orchiopexy. The Correct Answer is 1. Please review the study resources below ■ Teaching Points About 1 in 500 men with undescended testes develop testicular cancer, which occurs in the third to fourth decades of life. This patient is 17 years of age, and the testicle should be removed as it is not viable. One of the strongest arguments for early orchiopexy is to prevent testicular cancer. • The most common cancer is seminoma. It is treatable if diagnosed early. Therefore, urologists re orchiopexy. Young males can be taught a testicular exam, although this is not effective screen

#### Reading Assignment - Readings\_251001104920

Readings Due By: 10/7/2025




Airway Assessment

Access assigned articles by clicking on the Adjustiassignment, the on William Article." Mark Assignment as Complete when finished.

View Article

## Creating a Quiz

Go to Question Bank to build a quiz for self-study. You can access this in two places:



## **Quiz Options**

- Select from a wide range of question banks.
- Choose between "Study" and "Test" modes.
  - Study Mode: shows teaching points and feedback on incorrect answers.
  - Test Mode: answer all questions and review at the end.
- Select Difficulty Level and Age Range as desired.
- Narrow to specific topics.
- Choose number of questions for your quiz.
- Select "Create and Take Quiz" when done.
- Quiz will start immediately but will also appear in your created quizzes to complete at a later time.

## Track your progress

- You can see performance data such as correct and incorrect answers and a comparison to other users of the question bank.
- Pre and Post Tests based on your specialty allow you to compare your performance before and after.

#### **THANK YOU!**



Rowan Marye
Public Services Librarian
Nursing Liaison
rmarye@lsuhsc.edu
504-568-6100